


TSC Pd-D Fusion and Zeolite-Heat-Water

Frank Dodd (Tony) Smith, Jr - 2015
viXra 1502.0096

Abstract

TSC-Jitterbug Pd-D fusion with 147-atom Palladium clusters containing Deuterium in Sodium Zeolite Y cages produces energy that can be carried from the Deuterium electrons to the Palladium electrons to the Zeolite electrons, thus heating the Zeolite, which heat can be released as needed by reacting with D₂O to form steam. If the Water that is initially in the Zeolite and released as the Zeolite is heated by fusion is Hydrogen water, then the Hydrogen could contaminate the Pd cluster Deuterium and impair the TSC fusion process so all Zeolite used for TSC-Jitterbug Pd-D fusion should be of the form $A_mX_pO_{2p} \cdot nD_2O$ that is, all the water in the Zeolite used for Pd-D fusion should be Heavy Water D₂O which can be accomplished by taking ordinary Zeolite, then heating it to flush out all the Ordinary Water H₂O and then cooling it with Heavy Water D₂O to give $A_mX_pO_{2p} \cdot nD_2O$

TSC-Jitterbug Pd-D fusion with 147-atom Palladium clusters containing Deuterium



in Sodium Zeolite Y cages




is described in some detail in my papers at

<http://vixra.org/abs/1501.0234>

<http://vixra.org/abs/1502.0069>

If each 147-atom Palladium cluster is embedded into a Zeolite cage then the fusion energy can be carried from the Deuterium electrons to the Palladium electrons to the Zeolite electrons, thus heating the Zeolite, which heat can be released as needed by reacting with D2O to form steam.

“... Zeolite is a aluminosilicate mineral ... consist[ing] of three dimensional networks of AlO_4 and SiO_4 tetrahedra linked by ... oxygen atoms ... which contains crystal water. Its general chemical formula is




Where A represents Ca, Na, K, Ba, and Sr; X represents Al and Si. ...

When zeolites are heated, water molecules in it escape, and heat energy is stored in it ...[



]... in the meantime; when water molecules are adsorbed again, the heat energy in zeolites is released ...[




]...". (Han Baoqi, Yuan Hongyen, Yang Dequan, Liu Guoxi, Paper No. 9404 of Jilin Agricultural University, Changchun, Jilin Province, 130118, China, Utilization of natural zeolites for solar energy storage)

If the nH_2O that is initially in the Zeolite and released as the Zeolite is heated by fusion is Hydrogen water, then the Hydrogen could contaminate the Pd cluster Deuterium and impair the TSC fusion process

so

all Zeolite used for TSC-Jitterbug Pd-D fusion should be of the form



that is, all the water in the Zeolite used for Pd-D fusion should be Heavy Water D_2O which


can be accomplished by taking ordinary Zeolite

then heating it to flush out all the Ordinary Water D_2O

and then cooling it with Heavy Water D_2O to give $A_m X_p O_{2p} \cdot nD_2O$

so that


the overall process looks like



(Zeolite-Pd images adapted from Calvo and Carre in Nanotechnology 17 (2006) 1292-1299 and from <http://gwenbeads.blogspot.com/2014/04/infinite-skew-polyhedron-faujasite-4446.html>)

As to how the Water is configured in the Zeolite:

“... We report ... Monte Carlo simulations of water ... adsorption in [Zeolite] NaY ... faujasite ... The existence of cyclic water hexamers ... located in the 12-ring windows ...”



... recently disclosed by neutron diffraction experiments ... were ... observed in the case of NaY ...”. (Angela Di Lella, Nicolas Desbiens, Anne Boutin, Isabelle Demachy, Philippe Ungerer, Jean-Pierre Bellat, and Alain H. Fuchs, Phys. Chem. Chem. Phys. 8 (2006) 5396-5406)

According to a Fraunhofer press release at

www.fraunhofer.de/en/press/research-news/2014/march/sorption-energy-storage.html

“... Thermochemical heat storage systems based on a combination of zeolites and water have the potential to amass up to 180 kilowatt hours of energy per cubic meter ...”.